AN INFORMATION SHEET FOR SUPPORT PERSONS:

Type 1 diabetes in children and adolescents

what is hypoglycaemia?

Hypoglycaemia (low blood glucose level or hypo) occurs when the blood glucose level drops to less than 4mmol/L or when hypo symptoms are being experienced at a level close to 4mmol/L.

What are the main causes of a hypo?

- Delaying or missing meals or snacks
- · Being physically active

- Not eating enough carbohydrate
- · Having too much insulin

NOTE: Sometimes a hypo can occur for no obvious reason.

What are the symptoms of a hypo?

- A headache
- Trembling Looking pale
- Feeling hungry

- Sweating Crying

- Being irritable
 Feeling or acting confused

Treatment

Mild to moderate hypo

If the young person is conscious and has a blood glucose level less than 4 mmol/L, take the following steps.

Step 1

Give any one of the following. You may need to coax the young person to take it:

- 1/3–1/2 glass fruit juice or a small tetrapak
- 1/3–1/2 can soft drink (not Diet)
- 2–3 teaspoons honey or sugar
- Glucose tablets equivalent to 10–15 grams
- 5–7 jellybeans

Symptoms usually disappear after 10–15 minutes. However, if the blood glucose level remains low and symptoms are still present, repeat the treatment and stay with the young person.

Continued over...

If in doubt, do not waste time doing a blood glucose test. Blood glucose levels less than 4 mmol/L should be treated even when there are no symptoms.



what is hypoglycaemia?

Step 2

Depending on the child's insulin plan, whether or not they are on an insulin pump, their activity levels and how far away the next meal is, they may need additional carbohydrate such as fruit, a sandwich or biscuits to prevent their blood glucose level dropping again.

Severe hypo

If the young person has a fit or is unconscious, get emergency help fast!

- Do not attempt to give anything by mouth
- Lie the young person on their side in the recovery/coma position
- Call the ambulance (dial 000) OR give an injection of Glucagon* if it is available and you are trained to give it
- Stay with the young person until help arrives
- * Glucagon is a hormone that raises the blood glucose level and is injected into the muscle of the upper arm or the top of the leg. It is recommended to have Glucagon on hand in case of a severe hypo and be shown how to use it by a health professional.

References: Caring for Diabetes in Children and Adolescents – A Parent's Manual, Combined Children's Services of NSW, editors G Ambler, V Barron, C May and E Ambler; School Pack, International Diabetes Federation and Diabetes Australia.

Would you like to join Australia's leading diabetes organisation? > Dietary services > Free magazines > Children's services > Educational literature > Product discounts > Support groups For more information phone 1300 136 588 or visit your State/Territory Organisation's website: ACT www.diabetes-act.com.au NSW www.diabetesnsw.com.au

- **NT** www.healthylivingnt.org.au
- SA www.diabetessa.com.au
- VIC www.diabetesvic.org.au
- QLD www.diabetesqld.org.au
- TAS www.diabetestas.com.au
- WA www.diabeteswa.com.au

The design, content and production of this diabetes information sheet has been undertaken by:

- > Diabetes Australia NSW
- > Diabetes Australia Queensland
- > Diabetes ACT > Diabetes SA
- > Diabetes Australia Victoria
 > Diabetes Australia Tasmania
- Diabeles Australia -
 - > Diabetes WA > Healthy Living NT

The original medical and educational content of this information sheet has been reviewed by the Health Care and Education Committee of Diabetes Australia Ltd. Photocopying this publication in its original form is permitted for educational purposes only. Reproduction in any other form by third parties is prohibited. For any matters relating to this information sheet, please contact National Publications at dapubs@tpg.com.au or phone 02 9527 1951.

Health professionals: For bulk copies of this resource, contact Diabetes Australia in your State/Territory.